arXiv:1410.4020v1 [math.FA] 15 Oct 2014

STRONG CONVERGENCE THEOREMS BY A NEW HYBRID METHOD FOR EQUILIBRIUM PROBLEMS AND RELATIVELY NONEXPANSIVE MAPPINGS IN BANACH SPACES

SATTAR ALIZADEH[1] AND FRIDOUN MORADLOU[2]

Abstract. In this paper, we introduce a new modified Ishikawa iteration for finding a common element of the set of solutions of an equilibrium problem and the set of fixed points of relatively nonexpansive mappings in a Banach space. Our results generalize, extend and enrich some existing results in the literature.

1. Introduction

Throughout this paper, let E be a real Banach space with the dual space E^* and let C be a nonempty closed convex subset of E. Let f be a bifunction from $C \times C$ to $\mathbb{R}$. The equilibrium problem for $f : C \times C \to \mathbb{R}$ is to find $x \in E$ such that

$$f(x, y) \geq 0, \quad (y \in C). \tag{1.1}$$

The set of solutions of (1.1) is denoted by $EP(f)$, i.e.,

$$EP(f) = \{x \in C : \ f(x, y) \geq 0, \ \ \forall y \in C\}.$$

A self-mapping S of C is called nonexpansive if

$$\|Sx - Sy\| \leq \|x - y\|, \quad (x, y \in C).$$

We denote by $F(S)$ the set of fixed points of S.

Let $S : C \to E^*$ be a mapping and let $f(x, y) = \langle Sx, y - x\rangle$ for all $x, y \in C$. Then $z \in EP(f)$ if only if $\langle Sz, y - z\rangle \geq 0$ for all $y \in C$, i.e., z is a solution of the variational inequality $\langle Sx, y - x\rangle \geq 0$. So, the formulation (1.1) includes variational inequalities as special cases. Also, numerous problems in physics, optimization and economics reduce to find a solution of (1.1). Some methods have been proposed to solve the equilibrium problem; see for instance [5, 12, 15, 24, 20].

In the recent years, many authors studied the problem of finding a common element of the set of fixed points of a nonexpansive mapping and the set of solutions of an equilibrium problem in the framework of Hilbert spaces and Banach spaces, respectively; see for instance, [10, 11, 13, 17, 18, 26, 27, 30, 32] and the references therein.

2010 *Mathematics Subject Classification.* Primary 47H10, 47H09, 47J25, 47J05.

Key words and phrases. Equilibrium problems, Fixed point, Hybrid method, Strong convergence, Weak convergence.

Let C be a nonempty closed convex subset of a Banach space. In 1953, for a self-mapping S of C, Mann [21] defined the following iteration procedure:

$$\begin{cases} x_0 \in E \text{ chosen arbitrarily,} \\ x_{n+1} = \alpha_n x_n + (1-\alpha_n) S x_n, \end{cases} \tag{1.2}$$

where $0 \leq \alpha_n \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$,

Let K be a compact convex subset of a Hilbert space H. In 1974, for a Lipschitzian pseudocontractive self-mapping S of K, Ishikawa [16] defined the following iteration procedure:

$$\begin{cases} x_0 \in K \text{ chosen arbitrarily,} \\ y_n = \beta_n x_n + (1-\beta_n) S x_n, \\ x_{n+1} = \alpha_n x_n + (1-\alpha_n) S y_n, \end{cases} \tag{1.3}$$

where $0 \leq \beta_n \leq \alpha_n \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and he proved strong convergence of the sequence $\{x_n\}$ generated by the above iterative scheme if $\lim_{n\to\infty} \beta_n = 1$ and $\sum_{n=1}^{\infty}(1-\alpha_n)(1-\beta_n) = \infty$. By taking $\beta_n = 1$ for all $n \geq 0$ in (1.3), Ishikawa iteration process reduces to Mann iteration process.

In general, to gain the convergence in Mann and Ishikawa iteration processes, we must assume that underlying space E has elegant properties. For example, Reich [28] proved that if E is a uniformly convex Banach space with a Fréchet differentiable norm and if $\{\alpha_n\}$ is such that $\sum_{n=1}^{\infty} \alpha_n(1-\alpha_n) = \infty$, then the Mann iteration scheme converges weakly to a fixed point of T. However, we know that the Mann iteration process is weakly convergence even in a Hilbert space [14]. Also, Tan and Xu [36] proved that if E is a a uniformly convex Banach space which satisfies Opial's condition or whose norm is Fréchet differentiable and if $\{\alpha_n\}$ and $\{\beta_n\}$ are such that $\sum_{n=1}^{\infty} \alpha_n(1-\alpha_n)$ diverges, $\sum_{n=1}^{\infty} \alpha_n(1-\beta_n)$ converges and $\limsup \beta_n < 1$, then Ishikawa iteration process converges weakly to a fixed point of T.

It easy to see that process (1.3) is more general than the process (1.2). Also, for a Lipschitz pseudocontractive mapping in a Hilbert space, process (1.2) is not known to converge to a fixed point while the process (1.3) is convergence. In spite of these facts, researchers are interested to study the convergence theorems by process (1.2), because of the formulation of process (1.2) is simpler than that of (1.3) and if $\{\beta_n\}$ satisfies suitable conditions, we can gain a convergence theorem for process (1.3) on a convergence theorem for process (1.2).

In recent years, many authors have proved weak or strong convergence theorems for some nonlinear mappings by using various iteration processes in the framework of Hilbert spaces and Banach spaces, see, [3, 4, 25, 31, 35].

Recently, to obtain strong convergence, many mathematicians have been extensively considered modified processes. Nakajo and Takahashi [25] proposed the following modification of the Mann's iteration for a nonexpansive self-mapping S of a nonempty, closed convex

subset C in a Hilbert space H:

$$\begin{cases} x_0 \in C \text{ chosen arbitrarily}, \\ y_n = \alpha_n x_n + (1-\alpha_n) S x_n, \\ C_n = \{u \in C : \|y_n - u\| \leq \|x_n - u\|\}, \\ Q_n = \{u \in C : \langle x_n - u, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n} x_0, \end{cases}$$

where P_K denotes the metric projection from H onto a closed convex subset K of H. They proved strong convergence of the sequence $\{x_n\}$, if the sequence $\{\alpha_n\}$ is bounded above from one.

In 2005, Matsushita and Takahashi [23] introduced a new hybrid algorithim for a relatively self-mapping S of C in a Banach space E as follows:

$$\begin{cases} x_0 = x \in C \text{ chosen arbitrarily}, \\ y_n = J^{-1}(\alpha_n J x_n + (1-\alpha_n) J S x_n), \\ H_n = \{u \in C : \phi(u, y_n) \leq \phi(u, x_n)\}, \\ W_n = \{z \in C : \langle x_n - z, J x_n - J x \rangle \leq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n} x, n = 0, 1, 2, ..., \end{cases}$$

where J is the duality mapping on E and $\Pi_{F(S)}$ is the generalized projection from C onto $F(S)$. They proved that $\{x_n\}$ converges strongly to $\Pi_{F(S)} x$.

In 2006, Martinez-Yanes and Xu [22] introduced the following modified Ishikawa iteration process for a nonexpansive self-mapping S of a nonempty, closed convex subset C with $F(S) \neq \emptyset$ in a Hilbert space H:

$$\begin{cases} x_0 \in E \text{ chosen arbitrarily}, \\ z_n = \beta_n x_n + (1-\beta_n) S x_n \\ y_n = \alpha_n x_n + (1-\alpha_n) S z_n, \\ C_n = \{u \in C : \|y_n - u\|^2 \leq \|x_n - u\|^2 \\ \qquad\qquad +(1-\alpha_n)(\|z_n\|^2 - \|x_n\|^2 + 2\langle x_n - z_n, u \rangle \geq 0)\}, \\ Q_n = \{u \in C : \langle x_n - u, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n} x_0, \end{cases}$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$. They proved that if $\{\alpha_n\}$ bounded above from one and $\lim_{n\to\infty} \beta_n = 1$, then the sequence $\{x_n\}$ converges strongly to $P_{F(S)} x_0$.

In 2007, Tada and Takahashi [29] for finding an element of $EP(f) \cap F(S)$, introduced the following iterative scheme by using the hybrid projection method for a nonexpansive

self-mapping S of a nonempty, closed convex subset C in a Hilbert space:

$$\begin{cases} x_0 \in H \text{ chosen arbitrarily}, \\ u_n \in E \text{ such that } \quad f(u_n, y) + \frac{1}{r_n}\langle y - u_n, u_n - x_n\rangle \geq 0, \quad \forall\, y \in E \\ w_n = \alpha_n x_n + (1-\alpha_n) S u_n, \\ C_n = \{u \in H : \|w_n - u\| \leq \|x_n - u\|\}, \\ Q_n = \{u \in H : \langle x_n - u, x_n - x\rangle \leq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n} x, \end{cases}$$

for all $n \in \mathbb{N} \cup \{0\}$, where $\{\alpha_n\} \subset [a, b]$ for some $a, b \in (0, 1)$ and $\{r_n\} \subset (0, \infty)$ satisfies $\liminf_{n\to\infty} r_n > 0$. Thus, they proved that $\{x_n\}$ and $\{u_n\}$ converge strongly to u, where $u = P_{F(S)\cap EP(f)} x_0 \in F(S) \cap EP(f)$.

In 2009, Takahashi and Zembayashi [33] presented a new hybrid iterative method for finding an element of $EP(f) \cap F(S)$, by using the hybrid projection method for relatively nonexpansive self-mapping S of C in a Banach space E:

$$\begin{cases} x_0 \in C \text{ chosen arbitrarily}, \\ y_n = J^{-1}(\alpha_n J x_n + (1-\alpha_n) J S x_n), \\ u_n \in C \text{ such that } \quad f(u_n, y) + \frac{1}{r_n}\langle y - u_n, J u_n - J y_n\rangle \geq 0, \quad \forall\, y \in C \\ H_n = \{u \in C : \|w_n - u\| \leq \|x_n - u\|\}, \\ W_n = \{u \in C : \langle x_n - u, J x_n - J x\rangle \leq 0\}, \\ x_{n+1} = P_{H_n \cap W_n} x, \end{cases}$$

for all $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E, $\{\alpha_n\}$ is a sequence in $[0, 1]$ such that $\lim_{n\to\infty} \alpha_n(1 - \alpha_n) > 0$ and $\{r_n\}$ is a sequence in $[a, \infty)$ for some $a > 0$. They proved that the sequence $\{x_n\}$ converges strongly to $\Pi_{F(S)\cap EP(f)} x$, where $\Pi_{F(S)\cap EP(f)}$ is the generalized projection from C onto $F(S) \cap EP(f)$.

In this paper, employing the idea of Nakajo-Takahashi [25] and Takahashi-Zembayashi [33], we modify Ishikawa iteration process for finding a common element of the set of solution of an equilibrium problem and the set of fixed points of a relatively nonexpansive mapping by applying the hybrid projection method in a Banach space.

2. Preliminaries

Let E be a real Banach space with $\|.\|$ and dual space E^*. We denote the weak convergence and the strong convergence of $\{x_n\}$ to $x \in E$ by $x_n \rightharpoonup x$ and $x_n \to x$, respectively and denote by J the normalized duality mapping from E into 2^{E^*} defined by

$$Jx = \{x^* \in E^* : \ \langle x, x^*\rangle = \|x\|^2 = \|x^*\|^2\}$$

for all $x \in E$, where $\langle ., .\rangle$ denotes the generalized duality pairing between E and E^*. A Banach space E is said to be strictly convex if $\|\frac{x+y}{2}\| < 1$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $x \neq y$. It is also said to be uniformly convex if for every $\epsilon \in (0, 2]$, there exists a $\delta > 0$,

such that $\|\frac{x+y}{2}\| < 1-\delta$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $\|x-y\| \geq \epsilon$. Furthermore, E is called smooth if the limit

$$\lim_{t\to 0} \frac{\|x+ty\| - \|x\|}{t} \tag{2.1}$$

exists for all $x, y \in B_E = \{x \in E : \|x\| = 1\}$. It is also said to be uniformly smooth if the limit (2.1) is attained uniformly for all $x, y \in E$. A Banach space E is said to have the Kadec–Klee property if for every sequence $\{x_n\}$ in E that $x_n \rightharpoonup x$ and $\|x_n\| \to \|x\|$ then $x_n \to x$. It is known that if E is uniformly convex, then E has the Kadec–Klee property. Also, E is uniformly convex if and only if E^* is uniformly smooth. For more details see [1, 34].

Many properties of the normalized duality mapping J have been given in [1, 34]. We give some of those in the following:

(1) $J(0) = \{0\}$.
(2) For every $x \in E$, Jx is nonempty closed convex and bounded subset of E^*.
(3) If E^* is strictly convex, then J is single-valued.
(4) If E is strictly convex, then J is one-one, i.e., if $x \neq y$ then $Jx \cap Jy = \phi$.
(5) If E is reflexive, then J is onto.
(6) If E is smooth, then J is single-valued.
(7) if E is smooth and reflexive, then J is norm-to-weak continuous, that is, $Jx_n \rightharpoonup Jx$ whenever $x_n \to x$.
(8) If E is smooth, strictly convex and reflexive and $J^* : E^* \to 2^E$ is the normalized duality mapping on E^*, then $J^{-1} = J^*$, $JJ^* = I_{E^*}$ and $J^*J = I_E$, where I_E and I_{E^*} are the identity mapping on E and E^*, respectively.
(9) If E is uniformly convex and uniformly smooth, then J is uniformly norm-to-norm continuous on bounded sets of E and $J^{-1} = J^*$ is also uniformly norm-to-norm continuous on bounded sets of E^*.

Let C be a nonempty, closed convex subset of a smooth, strictly convex and reflexive Banach space E. We denote by ϕ the function $\phi : E \times E \to \mathbb{R}$ defined as follows:

$$\phi(x, y) = \|x\|^2 - \langle x, Jy\rangle + \|y\|^2,$$

for all $x, y \in E$. It is clear from the definition of the function ϕ that for all $x, y, z \in E$,

(1) $(\|y\| - \|x\|)^2 \leq \phi(x, y) \leq (\|y\| + \|x\|)^2$,
(2) $\phi(x, y) = \phi(x, z) + \phi(z, y) + 2\langle x - z, Jz - Jy\rangle$.

Observe that if E is in a Hilbert space then $\phi(x, y) = \|x - y\|^2$.

In 1996, Alber [2], defined the generalized projection mapping as follows:

Definition 2.1. Let C be a nonempty, closed convex subset of a smooth, strictly convex and reflexive Banach space E. The generalized projection $\Pi_C : E \to C$ is a mapping that assigns to an arbitrary point $x \in E$ the minimum point of the functional $\phi(x, y)$, i.e., $\Pi_C x = x_0$,

where x_0 is the solution to the minimization problem

$$\phi(x_0, x) = \inf_{y \in C} \phi(y, x).$$

Existence and uniqueness of the operator Π_C follows from the properties of the functional $\phi(x, y)$ and strict monotonicity of the mapping J.

Let S be a self-mapping of C. A point p in C is said to be an asymptotic fixed point of S [28], if there exists a sequence $\{x_n\}$ in C such that $x_n \rightharpoonup p$ and $\|x_n - Sx_n\| \to 0$. we denote by $\hat{F}(S)$ the set of all asymptotic fixed points of S. A self-mapping S of C is said to be relatively nonexpansive [5-7], if the following conditions are satisfied:

(1) $F(S)$ is nonempty;
(2) $\phi(u, Sx) \leq \phi(u, x)$, $\forall u \in F(S)$, $\forall x \in C$;
(3) $F(S) = \hat{F}(S)$.

Lemma 2.2. [23] *Let C be a nonempty, closed convex subset of a smooth, strictly convex and reflexive Banach space E, and let S be a relatively nonexpansive self-mapping of C. Then $F(S)$ is closed and convex.*

Lemma 2.3. [2] *Let C be a nonempty, closed convex subset of smooth, strictly convex and reflexive Banach space E. Then*

$$\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y),$$

for all $x \in C$ and all $y \in E$.

Lemma 2.4. [2] *Let C be a nonempty, closed convex subset of smooth, strictly convex and reflexive Banach space E, $x \in E$ and $z \in C$. Then*

$$z = \Pi_C x \Longleftrightarrow \langle y - z, Jx - Jz \rangle \leq 0,$$

for all $y \in C$.

Lemma 2.5. [19] *Let E be a smooth and uniformly convex Banach space and let $\{x_n\}$ and $\{y_n\}$ be sequences in E such that either $\{x_n\}$ or $\{y_n\}$ is bounded. If $lim_{n\to\infty}\phi(x_n, y_n) = 0$, then $lim_{n\to\infty}\|x_n - y_n\| = 0$.*

Lemma 2.6. [37] *Let E be a uniformly convex Banach space and $r > 0$. Then there exists a strictly increasing, continuous and convex function $g : [0, 2r] \to \mathbb{R}$ such that $g(0) = 0$ and*

$$\|tx + (1-t)y\|^2 \leq t\|x\|^2 + (1-t)\|y\|^2 - t(1-t)g(\|x - y\|),$$

for all $x, y \in B_r$ and $t \in [0, 1]$, where $B_r = \{z \in E : \|z\| \leq r\}$.

To study the equilibrium problem, for the bifunction $f : C \times C \to \mathbb{R}$, we assume that f satisfies the following conditions:

(A1) $f(x, x) = 0$ for all $x \in C$;
(A2) f is monotone, i.e., $f(x, y) + f(y, x) \leq 0$ for all $x, y \in C$;

(A3) for each $x, y, z \in C$,

$$\lim_{t \downarrow 0} f(tz + (1-t)x, y) \leq f(x, y);$$

(A4) for each $x \in C$, $y \mapsto f(x, y)$ is convex and lower semicontinuous.

The following lemma can be found in [5].

Lemma 2.7. *Let C be a nonempty, closed convex subset of a smooth, strictly convex and reflexive Banach space E, f be a bifunction from $C \times C$ to $\mathbb{R}$ satisfying $(A1) - (A4)$ and let $r > 0$ and $x \in E$. Then, there exists $z \in C$ such that*

$$f(z, y) + \frac{1}{r}\langle y - z, Jz - Jx\rangle \geq 0,$$

for all $y \in C$.

Lemma 2.8. [33] *Let C be a nonempty, closed convex subset of a smooth, strictly convex and reflexive Banach space E, f be a bifunction from $C \times C$ to $\mathbb{R}$ satisfying $(A1) - (A4)$ and let $r > 0$ and $x \in E$. Define a mapping $T_r : E \to C$ as follows:*

$$T_r(x) = \{z \in C : f(z, y) + \frac{1}{r}\langle y - z, Jz - Jx\rangle \geq 0, \ \forall y \in C\},$$

for all $x \in E$. Then, the following statements hold:

(i) *T_r is singel-valued;*
(ii) *T_r is firmly nonexpansive-type, i.e., for all $x, y \in E$,*

$$\langle T_r x - T_r y, JT_r x - JT_r y\rangle \leq \langle T_r x - T_r y, Jx - Jy\rangle;$$

(iii) *$F(T_r) = EP(f)$;*
(iv) *EP(f) is closed and convex and T_r is relatively nonexpansive mapping.*

3. Main Results

In this section, we prove the strong convergence theorem for finding a common element of the set of solution of an equilibrium problem and the set of fixed points of a relatively nonexpansive mapping.

Theorem 3.1. *Let C be a nonempty closed convex subset of uniformly smooth and uniformly convex Banach space E. Let f be a bifunction from $C \times C$ to $\mathbb{R}$ satisfying $(A1) - (A4)$ and S be a relatively nonexpansive self-mapping of C with $F(S) \cap EP(f) \neq \phi$. Assume that $0 < a \leq \alpha_n \leq 1$ and $\{r_n\} \subset (0, \infty)$ satisfies $\liminf_{n\to\infty} r_n > 0$ and $\{\beta_n\}$ is sequence in $[0, 1]$ such that $\liminf_{n\to\infty} \beta_n(1-\beta_n) > 0$. If $\{x_n\}$ and $\{u_n\}$ be sequences generated by $x = x_1 \in C$*

and

$$\begin{cases} z_n = J^{-1}(\beta_n Jx_n + (1-\beta_n)JSx_n), \\ u_n \in E \text{ such that } \quad f(u_n, y) + \frac{1}{r_n}\langle y - u_n, Ju_n - Jz_n\rangle \geq 0, \quad \forall\, y \in E \\ y_n = J^{-1}(\alpha_n Jz_n + (1-\alpha_n)Ju_n), \\ C_n = \{v \in C : \phi(v, y_n) \leq \phi(v, x_n)\}, \\ Q_n = \{z \in C : \langle x_n - z, Jx_n - Jx\rangle \leq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n} x, \end{cases}$$

for all $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to $\Pi_{F(S)\cap EP(f)}x$, where $\Pi_{F(S)\cap EP(f)}$ is the generalized projection of E onto $F(S) \cap EP(f)$.

Proof. First, we show that $C_n \cap Q_n$ is closed and convex. It is easily seen that C_n is closed and Q_n is closed and convex for all $n \in \mathbb{N}$. Since

$$\phi(v, y_n) \leq \phi(v, x_n) \Longleftrightarrow \|y_n\|^2 - \|x_n\|^2 - 2\langle v, Jy_n - Jx_n\rangle \geq 0,$$

for all $v \in C_n$. Then $v \in C_n$ is convex. So $C_n \cap Q_n$ is closed and convex for all $n \in \mathbb{N}$. Let $u \in F(S) \cap EP(f)$. From $u_n = T_{r_n} z_n$ and using Lemma 2.8(iv), we can conclude that T_{r_n} are relatively nonexpansive, in addition, S is relatively nonexpansive, so by the convexity of $\|.\|^2$ we get

$$\begin{aligned} \phi(u, u_n) &= \phi(u, T_{r_n} z_n) \\ &\leq \phi(u, z_n) \\ &= \phi(u, J^{-1}(\beta_n Jx_n + (1-\beta_n)JSx_n) \\ &= \|u\|^2 - 2\langle u, \beta_n Jx_n + (1-\beta_n)JSx_n\rangle + \|\beta_n Jx_n + (1-\beta_n)JSx_n\|^2 \\ &\leq \|u\|^2 - 2\beta_n\langle u, Jx_n\rangle - 2(1-\beta_n)\langle u, JSx_n\rangle + \beta_n\|x_n\|^2 + (1-\beta_n)\|Sx_n\|^2 \\ &= \beta_n\phi(u, x_n) + (1-\beta_n)\phi(u, Sx_n) \\ &\leq \phi(u, x_n), \end{aligned} \tag{3.1}$$

and therefore

$$\begin{aligned} \phi(u, y_n) &= \phi(u, J^{-1}(\alpha_n Jz_n + (1-\alpha_n)Ju_n) \\ &= \|u\|^2 - 2\langle u, \alpha_n Jz_n + (1-\alpha_n)Ju_n\rangle + \|\alpha_n Jz_n + (1-\alpha_n)Ju_n\|^2 \\ &\leq \|u\|^2 - 2\alpha_n\langle u, Jz_n\rangle - 2(1-\alpha_n)\langle u, Ju_n\rangle + \alpha_n\|z_n\|^2 + (1-\alpha_n)\|u_n\|^2 \\ &= \alpha_n\phi(u, z_n) + (1-\alpha_n)\phi(u, u_n) \\ &\leq \alpha_n\phi(u, x_n) + (1-\alpha_n)\phi(u, x_n) \\ &= \phi(u, x_n), \end{aligned} \tag{3.2}$$

So $u \in C_n$ for all $n \in \mathbb{N}$. Hence,

$$F(S) \cap EP(f) \subset C_n.$$

Now, using induction, we will show that $F(S) \cap EP(f) \subset C_n \cap Q_n$ for all $n \in \mathbb{N}$. For $n = 1$, we have $x_1 = x \in C$ and $Q_1 = C$, therefore $F(S) \cap EP(f) \subset Q_1$ and hence

$F(S) \cap EP(f) \subset C_1 \cap Q_1$. Assume that $F(S) \cap EP(f) \subset C_n \cap Q_n$ for some n. Since $x_{n+1} = \Pi_{C_n \cap Q_n} x$, by Lemma 2.4, we have

$$\langle x_{n+1} - z, Jx - Jx_{n+1} \rangle \geqslant 0, \tag{3.3}$$

for all $z \in C_n \cap Q_n$. By the induction assumption, we have $F(S) \cap EP(f) \subset C_n \cap Q_n$, so we can conclude that for all $u \in F(S) \cap EP(f)$ the inequality (3.3) holds. On the other hand, by the definition of Q_{n+1}, we obtain $F(S) \cap EP(f) \subset Q_{n+1}$. Therefore, $F(S) \cap EP(f) \subset C_{n+1} \cap Q_{n+1}$. So $\{x_n\}$ is well-defined.

Notice that the definition of Q_n implies $x_n = \Pi_{Q_n} x$. By Lemma 2.3 we get

$$\phi(x_n, x) = \phi(\Pi_{Q_n} x, x) \leq \phi(u, x) - \phi(u, \Pi_{Q_n} x) \leq \phi(u, x), \tag{3.4}$$

for all $u \in F(S) \cap EP(f) \subset Q_n$. This yields that $\{\phi(x_n, x)\}$ is bounded. Therefore $\{x_n\}$, $\{Sx_n\}$, $\{y_n\}$ and $\{z_n\}$ are bounded.

Since $x_{n+1} = \Pi_{C_n \cap Q_n} x \in C_n \cap Q_n \subset Q_n$, and $x_n = \Pi_{Q_n} x$, from the definition of Π_{Q_n} we have

$$\phi(x_n, x) \leq \phi(x_{n+1}, x), \tag{3.5}$$

for all $n \in \mathbb{N}$. It follows from (3.4) and (3.5), the sequence $\{\phi(x_n, x)\}$ is bounded and nondecreasing . So, $\lim_{n\to\infty}\{\phi(x_n, x)\}$ exists. by using Lemma 2.3 and $x_n = \Pi_{Q_n} x$, we also get

$$\begin{aligned} \phi(x_{n+1}, x_n) &= \phi(x_{n+1}, \Pi_{Q_n} x) \\ &\leq \phi(x_{n+1}, x) - \phi(\Pi_{Q_n} x, x) \\ &= \phi(x_{n+1}, x) - \phi(x_n, x), \end{aligned} \tag{3.6}$$

for all $n \in \mathbb{N}$. This yields that

$$\lim_{n\to\infty} \phi(x_{n+1}, x_n) = 0. \tag{3.7}$$

Since $x_{n+1} = \Pi_{C_n \cap Q_n} x \in C_n$, we have

$$\phi(x_{n+1}, y_n) \leq \phi(x_{n+1}, x_n), \tag{3.8}$$

for all $n \in \mathbb{N}$. from (3.7) and (3.8) we have

$$\lim_{n\to\infty} \phi(x_{n+1}, y_n) = 0. \tag{3.9}$$

From (3.7) and (3.9), since E is uniformly convex and smooth, from Lemma 2.5 we get

$$\lim_{n\to\infty} \|x_{n+1} - x_n\| = \lim_{n\to\infty} \|x_{n+1} - y_n\| = 0.$$

So, we obtain

$$\lim_{n\to\infty} \|x_n - y_n\| = 0. \tag{3.10}$$

Since J is uniformly norm-to-norm continuous on bounded sets, from (3.10) we get

$$\lim_{n\to\infty} \|Jx_n - Jy_n\| = 0$$

Let $r = \sup_{n\in\mathbb{N}}\{\|x_n\|, \|Sx_n\|\}$. Since E is a uniformly smooth Banach space, we can conclude that E^* is a uniformly convex Banach space. So, by Lemma 2.6 there exists a continuous,

strictly increasing and convex function $g : [0, 2r] \to \mathbb{R}$ whit $g(0) = 0$ and for $u \in F(S) \cap EP(f)$ we have

$$\begin{aligned}
\phi(u, z_n) &= \phi(u, J^{-1}(\beta_n Jx_n + (1-\beta_n)JSx_n) \\
&= \|u\|^2 - 2\langle u, \beta_n Jx_n + (1-\beta_n)JSx_n\rangle + \|\beta_n Jx_n + (1-\beta_n)JSx_n\|^2 \\
&\leq \|u\|^2 - 2\beta_n\langle u, Jz_n\rangle - 2(1-\beta_n)\langle u, JSu_n\rangle + \beta_n\|z_n\|^2 + (1-\beta_n)\|Su_n\|^2 \\
&\qquad - \beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|) \\
&= \beta_n\phi(u, x_n) + (1-\beta_n)\phi(u, Sx_n) - \beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|) \\
&\leq \phi(u, x_n) - \beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|),
\end{aligned} \tag{3.11}$$

and therefore

$$\begin{aligned}
\phi(u, y_n) &= \phi(u, J^{-1}(\alpha_n Jz_n + (1-\alpha_n)Ju_n) \\
&\leq \alpha_n\phi(u, z_n) + (1-\alpha_n)\phi(u, x_n) \\
&\leq \alpha_n\phi(u, x_n) - \alpha_n\beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|) + (1-\alpha_n)\phi(u, x_n) \\
&\leq \phi(u, x_n) - \alpha_n\beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|),
\end{aligned} \tag{3.12}$$

hence

$$\alpha_n\beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|) \leq \phi(u, x_n) - \phi(u, y_n).$$

Since $0 < a \leq \alpha_n \leq 1$, it is easy to see that

$$a\beta_n(1-\beta_n)g(\|Jx_n - JSx_n\|) \leq \phi(u, x_n) - \phi(u, y_n). \tag{3.13}$$

On the other hand, we have

$$\begin{aligned}
\phi(u, x_n) - \phi(u, y_n) &= \|x_n\|^2 - \|y_n\|^2 - 2\langle u, Jx_n - Jy_n\rangle \\
&\leq |\|x_n\|^2 - \|y_n\|^2| + 2|\langle u, Jx_n - Jy_n\rangle| \\
&\leq |\|x_n\| - \|y_n\||(\|x_n\| + \|y_n\|) + 2\|u\|\|Jx_n - Jy_n\| \\
&\leq |\|x_n - y_n\||(\|x_n\| + \|y_n\|) + 2\|u\|\|Jx_n - Jy_n\|,
\end{aligned} \tag{3.14}$$

so

$$\lim_{n\to\infty}(\phi(u, x_n) - \phi(u, y_n)) = 0 \tag{3.15}$$

Since $\liminf_{n\to\infty} \beta_n(1-\beta_n) > 0$, it follows from (3.13) that

$$\lim_{n\to\infty} g(\|Jx_n - JSx_n\|) = 0.$$

Hence, from property of g we get

$$\lim_{n\to\infty} \|JSx_n - Jx_n\| = 0. \tag{3.16}$$

Since J^{-1} is also uniformly norm-to-norm continuous on bounded sets, we have

$$\lim_{n\to\infty} \|x_n - Sx_n\| = 0. \tag{3.17}$$

Notice that $Jz_n - Jx_n = (1-\beta_n)(JSx_n - Jx_n)$, from (3.16) we can conclude that

$$\lim_{n\to\infty} \|Jz_n - Jx_n\| = \lim_{n\to\infty}(1-\beta_n)\|JSx_n - Jx_n\| = 0.$$

Since J^{-1} is also uniformly norm-to-norm continuous on bounded sets, we obtain

$$\lim_{n\to\infty} \|z_n - x_n\| = 0. \tag{3.18}$$

Since

$$\|y_n - z_n\| \leq \|y_n - x_n\| + \|x_n - z_n\|,$$

from (3.10), (3.18) and last inequality we have

$$\lim_{n\to\infty} \|y_n - z_n\| = 0. \tag{3.19}$$

Since J is uniformly norm-to-norm continuous on bounded sets from (3.19), we get

$$\lim_{n\to\infty} \|Jy_n - Jz_n\| = 0. \tag{3.20}$$

Noticing that $Jy_n = \alpha_n Jz_n + (1-\alpha_n)Ju_n$ and $\alpha_n < b < 1$, we obtain

$$\|Jz_n - Ju_n\| = \frac{1}{1-\alpha_n}\|Jy_n - Jz_n\| \leq \frac{1}{1-b}\|Jy_n - Jz_n\|.$$

From (3.20) and last inequality we get

$$\lim_{n\to\infty} \|Jz_n - Ju_n\| = 0. \tag{3.21}$$

Since J^{-1} is also uniformly norm-to-norm continuous on bounded sets, we get

$$\lim_{n\to\infty} \|z_n - u_n\| = 0. \tag{3.22}$$

Since $\{x_n\}$ is bounded, there is a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} \rightharpoonup \hat{x}$.

Since S is relatively nonexpansive, we can conclude from (3.17) that $\hat{x} \in \hat{F}(S) = F(S)$. We show that $\hat{x} \in EP(f)$. From $x_{n_k} \rightharpoonup \hat{x}$, (3.18) and (3.22), we have $z_{n_k} \rightharpoonup \hat{x}$ and $u_{n_k} \rightharpoonup \hat{x}$.

Since $\liminf_{n\to\infty} r_n > 0$, from (3.21), we derive that

$$\lim_{n\to\infty} \left\| \frac{Ju_n - Jz_n}{r_n} \right\| = \lim_{n\to\infty} \frac{1}{r_n}\|Ju_n - Jz_n\| = 0. \tag{3.23}$$

Since $u_n = T_{r_n} z_n$, we get

$$f(u_n, z) + \frac{1}{r_n}\langle z - u_n, Ju_n - Jz_n\rangle \geq 0, \tag{3.24}$$

for all $y \in C$. Substituting n by n_k in (3.24) and by using condition $(A2)$, we obtain

$$\frac{1}{r_{n_k}}\langle z - u_{n_k}, Ju_{n_k} - Jz_{n_k}\rangle \geq -f(u_{n_k}, y) \geq f(y, u_{n_k}),$$

for all $y \in C$. Letting $k \to \infty$, it follows from (3.23) and condition $(A4)$ that

$$0 \geq f(y, \hat{x}),$$

for all $y \in C$. Suppose that $t \in (0, 1]$, $y \in C$ and $y_t = ty + (1-t)\hat{x}$. Therefore, $y_t \in C$ and so $f(y_t, \hat{x}) \leq 0$. Hence

$$0 = f(y_t, y_t) \leq tf(y_t, y) + (1-t)f(y_t, \hat{x}) \leq tf(y_t, y),$$

and dividing by t, we have $f(y_t, y) \geq 0$, for all $y \in E$. By taking the limit as $t \downarrow 0$ and using $(A3)$, we get $\hat{x} \in EP(f)$.

put $\omega = \Pi_{F(S)\cap EP(f)}x$. From $x_{n+1} = \Pi_{C_n\cap Q_n}x$ and $\omega \in F(S)\cap EP(f) \subset C_n \cap Q_n$, we have

$$\phi(x_{n+1}, x) \leq \phi(\omega, x).$$

Since $\|.\|$ is weakly lower semicontinuous, we get

$$\begin{aligned}\phi(\hat{x}, x) =& \|\hat{x}\|^2 - \langle \hat{x}, Jx\rangle + \|x\|^2 \\ &\leq \liminf_{n\to\infty}(\|x_{n_k}\|^2 - \langle x_{n_k}, Jx\rangle + \|x\|^2) \\ &= \liminf_{n\to\infty}\phi(x_{n_k}, x) \\ &\leq \limsup_{n\to\infty}\phi(x_{n_k}, x) \\ &\leq \phi(\omega, x).\end{aligned}$$

By the definition of $\Pi_{F(S)\cap EP(f)}$, we get $\hat{x} = \omega$, so we get $\lim_{n\to\infty}\phi(x_{n_k}, x) = \phi(\omega, x)$. Hence we obtain

$$\begin{aligned}0 &= \lim_{n\to\infty}(\phi(x_{n_k}, x) - \phi(\omega, x)) \\ &= \lim_{n\to\infty}(\|x_{n_k}\|^2 - \|\omega\|^2 - 2\langle x_{n_k} - \omega, Jx\rangle) \\ &= \lim_{n\to\infty}(\|x_{n_k}\|^2 - \|\omega\|^2).\end{aligned}$$

Since E has Kadac–Klee property, we can conclude that $x_{n_k} \to \omega = \Pi_{F(S)\cap EP(f)}x$. Hence $\{x_n\}$ converges strongly to $\Pi_{F(S)\cap EP(f)}x$ □

Corollary 3.2. *Let C be a nonempty, closed convex subset of uniformly smooth and uniformly convex Banach space E and S be a relatively nonexpansive self-mapping of C with $F(S) \neq \phi$. Assume that $0 < a \leq \alpha_n \leq 1$ and $\{\beta_n\}$ is sequence in $[0, 1]$ such that $\liminf_{n\to\infty}\beta_n(1-\beta_n) > 0$. If $\{x_n\}$ and $\{u_n\}$ be sequences generated by $x = x_1 \in C$ and*

$$\begin{cases} z_n = J^{-1}(\beta_n Jx_n + (1-\beta_n)JSx_n), \\ u_n \in E \text{ such that } \quad \langle y - u_n, Ju_n - Jz_n\rangle \geq 0, \quad \forall\, y \in E \\ y_n = J^{-1}(\alpha_n Jz_n + (1-\alpha_n)Ju_n), \\ C_n = \{v \in C : \phi(v, y_n) \leq \phi(v, x_n)\}, \\ Q_n = \{z \in C : \langle x_n - z, Jx_n - Jx\rangle \leq 0\}, \\ x_{n+1} = \Pi_{C_n\cap Q_n}x, \end{cases}$$

for all $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to $\Pi_{F(S)}x$, where $\Pi_{F(S)}$ is the generalized projection of E onto $F(S)$.

Proof. Letting $f(x, y) = 0$ for all $x, y \in E$ and $r_n = 1$ for all $n \in \mathbb{N}$, in Theorem 3.1, we get the desired result. □

Corollary 3.3. *Let C be a nonempty, closed convex subset of uniformly smooth and uniformly convex Banach space E and f be a bifunction from $C \times C$ to $\mathbb{R}$ satisfying $(A1)-(A4)$ and S be a relatively nonexpansive self-mapping of C with $F(S)\cap EP(f) \neq \phi$. Assume*

that $\{r_n\} \subset (0,\infty)$ *satisfies* $\liminf_{n\to\infty} r_n > 0$ *and* $\{\beta_n\}$ *is sequence in* $[0,1]$ *such that* $\liminf_{n\to\infty} \beta_n(1-\beta_n) > 0$. *If* $\{x_n\}$ *and* $\{u_n\}$ *be sequences generated by* $x = x_1 \in C$ *and*

$$\begin{cases} z_n = J^{-1}(\beta_n Jx_n + (1-\beta_n)JSx_n), \\ u_n \in E \text{ such that } \quad f(u_n, y) + \frac{1}{r_n}\langle y - u_n, Ju_n - Jz_n\rangle \geq 0, \quad \forall\, y \in E \\ C_n = \{v \in C : \phi(v, z_n) \leq \phi(v, x_n)\}, \\ Q_n = \{z \in C : \langle x_n - z, Jx_n - Jx\rangle \leq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n} x, \end{cases}$$

for all $n \in \mathbb{N}$. *Then,* $\{x_n\}$ *converges strongly to* $\Pi_{F(S)\cap EP(f)}x$, *where* $\Pi_{F(S)\cap EP(f)}$ *is the generalized projection of* E *onto* $F(S) \cap EP(f)$.

Proof. Letting $\alpha_n = 1$, in Theorem 3.1, we get the desired result. □

Corollary 3.4. *Let* C *be a nonempty closed convex subset of uniformly smooth and uniformly convex Banach space* E *and* f *be a bifunction from* $C \times C$ *to* $\mathbb{R}$ *satisfying* $(A1) - (A4)$ *with* $EP(f) \neq \phi$. *Assume that* $0 < a \leq \alpha_n \leq 1$ *and* $\{r_n\} \subset (0,\infty)$ *satisfies* $\liminf_{n\to\infty} r_n > 0$ *and* $\{\beta_n\}$ *is sequence in* $[0,1]$ *such that* $\liminf_{n\to\infty} \beta_n(1-\beta_n) > 0$. *If* $\{x_n\}$ *and* $\{u_n\}$ *be sequences generated by* $x = x_1 \in C$ *and*

$$\begin{cases} u_n \in E \text{ such that } \quad f(u_n, y) + \frac{1}{r_n}\langle y - u_n, Ju_n - Jx_n\rangle \geq 0, \quad \forall\, y \in E \\ y_n = J^{-1}(\alpha_n Jx_n + (1-\alpha_n)Ju_n), \\ C_n = \{v \in C : \phi(v, y_n) \leq \phi(v, x_n)\}, \\ Q_n = \{z \in C : \langle x_n - z, Jx_n - Jx\rangle \leq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n} x, \end{cases}$$

for all $n \in \mathbb{N}$. *Then,* $\{x_n\}$ *converges strongly to* $\Pi_{EP(f)}x$, *where* $\Pi_{EP(f)}$ *is the generalized projection of* E *onto* $EP(f)$.

Proof. Letting $S = I$ in Theorem 3.1, we get the desired result. □

References

[1] R. P. Agarwal, Donal O'Regan and D. R. Sahu, *Fixed Point Theory for Lipschitzian-type Mappings with Applications*, Springer 2009.

[2] Ya. I. Alber, *Metric and generalized projection operators in Banach spaces: Properties and applications,* in: A. G. Kartsatos (Ed.), Theory and Applications of Nonlinear Operators of Accretive and Monotone Type, Marcel Dekker, New York, (1996), pp. 15–50.

[3] S. Alizadeh and F. Moradlou, *Strong convergence theorems for m-generalized hybrid mappings in Hilbert spaces,* to appear in Topol. Methods Nonlinear Anal.

[4] S. Alizadeh and F. Moradlou, *A strong convergence theorem for equilibrium problems and generalized hybrid mappings,* Mediterr. J. Math. DOI 10.1007/s00009-014-0462-6.

[5] E. Blum and W. Oettli, *From optimization and variational inequalities to equilibrium problems, Mathematics Students,* **63** (1994), 123–145.

[6] D. Butnariu, S. Reich and A. J. Zaslavski, *Asymptotic behavior of relatively nonexpansive operators in Banach spaces,* J. Appl. Anal. **7** (2001) 151–174.

[7] D. Butnariu, S. Reich and A. J. Zaslavski, *Weak convergence of orbits of nonlinear operators in reflexive Banach spaces,* Numer. Funct. Anal. Optim. **24** (2003) 489–508.

[8] Y. Censor and S. Reich, *Iterations of paracontractions and firmly nonexpansive operators with applications to feasibility and optimization,* Optimization **37** (1996) 323–339.

[9] C. E. Chidume and S. A. Mutangadura, *An example on the Mann iteration method for Lipschitz pseudocontractions,* Proc. Am. Math. Soc. **129** (2001) 2359-2363.
[10] L. C. Ceng and J. C. Yao, *A hybrid iterative scheme for mixed equilibrium problems and fixed point problems,* J. Comput. Appl. Math. **214** (2008) 186–201.
[11] L. C. Ceng, S. Al-Homidan and Q. H. Ansari, J. C. Yao, *An iterative scheme for equilibrium problems and fixed point problems of strict pseudo-contraction mappings,* J. Comput. Appl. Math. **223** (2009) 967–974.
[12] P. L. Combettes and S. A. Hirstoaga, *Equilibrium programming in Hilbert spaces,* J. Nonlinear Convex Anal. **6** (2005) 117–136.
[13] S. D. Flam and A. S. Antipin, *Equilibrium progamming using proximal-link algolithms,* Math. Program. **78** (1997) 29–41.
[14] A. Genel and J. Lindenstrass, *An example concerning fixed points,* Israel J. Math. **22** (1975) 81-86.
[15] F. Giannessi, A. Maugeri and P. M. Pardalos, *Equilibrium Problems: Nonsmooth Optimization and Variational Inequality Models,* Kluwer Academics Publishers, Dordrecht, Holland, 2001.
[16] S. Ishikawa, *Fixed points by a new iteration method*, Proc. Amer. Math. Soc. **40** (1974) 147–150.
[17] C. Jaiboon and P. Kumam, *A hybrid extragradient viscosity approximation method for solving equilibrium problems and fixed point problems of infinitely many nonexpansive mappings,* Fixed Point Theory Appl. (2009) Article ID 374815, 32 pages.
[18] C. Jaiboon and P. Kumam, H. W. Humphries, *Weak convergence theorem by an extragradient method for variational inequality, equilibrium and fixed point problems,* Bull. Malays. Math. Sci. Soc. (2) **32** (2) (2009) 173–185.
[19] S. Kamimura and W. Takahashi, *Strong convergence of a proximal-type algorithm in a Banach space,* SIAM J. Optim. **13** (2002) 938–945.
[20] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and their Applications,* Academic Press, New York, 1980.
[21] W. R. Mann, *Mean value methods in iteration,* Proc. Amer. Math. Soc. **4** (1953) 506–510.
[22] C. Martinez-Yanes and H. K. Xu, *Strong convergence of CQ method fixed point biteration processes,* Nonlinear Anal. **64** (2006) 2400–2411.
[23] S. Matsushita and W. Takahashi, *A strong convergence for relatively nonexpansive mappings in a Banach space,* J. Approx. Theory **134** (2005) 257–266.
[24] A. Moudafi and M. Thera, *Proximal and dynamical approaches to equilibrium problems, in: Lecture Note in Economics and Mathematical Systems,* vol. **477**, Springer-Verlag, New York, (1999), pp. 187–201.
[25] K. Nakajo and W. Takahashi, *Strong convergence theorems for nonexpansive mapping and nonexpansive mappings*, J. Math. Anal. Appl. **279** (2003) 372–379.
[26] S. Plubtieng and P. Kumam, *Weak convergence theorem for monotone mappings and a countable family of nonexpansive mappings,* J. Comput. Appl. Math. **224** (2009) 614–621.
[27] X. Qin, Y. J. Cho and S. M. Kang, *Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces,* J. Comput. Appl. Math. **225** (2009) 20–30.
[28] S. Reich, *A weak convergence theorem for the alternating method with Bregman distance,* in: A.G. Kartsatos (Ed.), Theory and Applications of Nonlinear Operators of Accretive and Monotone Type, Marcel Dekker, New York, (1996), pp. 313–318.
[29] A. Tada and W. Takahashi, *Weak and strong convergence theorems for a nonexpansive mappings and an equilibrium problem,* J. Optim. Theory Appl. **133** (2007) 359–370.
[30] S. Takahashi and W. Takahashi, *Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces,* J. Math. Anal. Appl. **331** (2007) 506–515.
[31] W. Takahashi and M. Toyoda, *Weak convergence theorems for nonexpansive mappings and monotone mappings,* J. Optim. Theory Appl. **118** (2003) 417-428.
[32] W. Takahashi and K. Zembayashi, *Strong convergence theorems by hybrid methods for equilibrium problems and relatively nonexpansive mapping,* Fixed Point Theory Appl. (2008) Article ID 528476, 11 pages.
[33] W. Takahashi, K. Zembayashi, *Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces,* Nonlinear Anal. **70** (2009) 45–57.
[34] W. Takahashi, *Introduction to Nonlinear and Convex Analysis*, Yokohoma Publishers, Yokohoma, 2009.
[35] W. Takahashi and J. -C. Yao, *weak convergence theorems for generalized hybrid mappings in Banach spaces,* Journal of Nonlinear Analysis and Optimazition. **2** (2011) No. 1, 147–158.
[36] K. K. Tan, H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. **178** (2) (1993) 301–308.

[37] H. K. Xu, *Inequalities in Banach spaces with application,* Nonlinear Anal. **16** (1991) 1127–1138.

[1,2]Department of Mathematics
Sahand University of Technology
Tabriz, Iran
E-mail address: [1] sa_alizadeh@sut.ac.ir
E-mail address: [2] moradlou@sut.ac.ir & fridoun.moradlou@gmail.com